

Quotient Isomorphism Invariants of a Finitely Generated Coxeter Group

Michael Mihalik, John Ratcliffe, and Steven Tschantz
Mathematics Department, Vanderbilt University,
Nashville TN 37240, USA

1 Introduction

The isomorphism problem for finitely generated Coxeter groups is the problem of deciding if two finite Coxeter matrices define isomorphic Coxeter groups. Coxeter [4] solved this problem for finite irreducible Coxeter groups. Recently there has been considerable interest and activity on the isomorphism problem for arbitrary finitely generated Coxeter groups.

In this paper we describe a family of isomorphism invariants of a finitely generated Coxeter group W . Each of these invariants is the isomorphism type of a quotient group W/N of W by a characteristic subgroup N . The virtue of these invariants is that W/N is also a Coxeter group. For some of these invariants, the isomorphism problem of W/N is solved and so we obtain isomorphism invariants that can be effectively used to distinguish isomorphism types of finitely generated Coxeter groups.

We emphasize that even if the isomorphism problem for finitely generated Coxeter groups is eventually solved, several of the algorithms described in our paper will still be useful because they are computational fast and would most likely be incorporated into an efficient computer program that determines if two finite rank Coxeter systems have isomorphic groups.

In §2, we establish notation. In §3, we describe two elementary quotienting operations on a Coxeter system that yields another Coxeter system. In §4, we describe the even part isomorphism invariant of a finitely generated Coxeter group. In §5, we review some matching theorems. In §6, we describe the even isomorphism invariant of a finitely generated Coxeter group. In §7, we define basic characteristic subgroups of a finitely generated Coxeter

group. In §8, we describe the spherical rank two isomorphism invariant of a finitely generated Coxeter group. In §9, we make some concluding remarks.

2 Preliminaries

A *Coxeter matrix* is a symmetric matrix $M = (m(s, t))_{s, t \in S}$ with $m(s, t)$ either a positive integer or infinity and $m(s, t) = 1$ if and only if $s = t$. A *Coxeter system* with Coxeter matrix $M = (m(s, t))_{s, t \in S}$ is a pair (W, S) consisting of a group W and a set of generators S for W such that W has the presentation

$$W = \langle S \mid (st)^{m(s, t)} : s, t \in S \text{ and } m(s, t) < \infty \rangle.$$

We call the above presentation of W , the *Coxeter presentation* of (W, S) . If (W, S) is a Coxeter system with Coxeter matrix $M = (m(s, t))_{s, t \in S}$, then the order of st is $m(s, t)$ for each s, t in S by Prop. 4, p. 92 of Bourbaki [3], and so a Coxeter system (W, S) determines its Coxeter matrix; moreover, any Coxeter matrix $M = (m(s, t))_{s, t \in S}$ determines a Coxeter system (W, S) where W is defined by the corresponding Coxeter presentation. If (W, S) is a Coxeter system, then W is called a *Coxeter group* and S is called a set of *Coxeter generators* for W , and the cardinality of S is called the *rank* of (W, S) . A Coxeter system (W, S) has finite rank if and only if W is finitely generated by Theorem 2 (iii), p. 20 of Bourbaki [3].

Let (W, S) be a Coxeter system. A *visual subgroup* of (W, S) is a subgroup of W of the form $\langle A \rangle$ for some $A \subset S$. If $\langle A \rangle$ is a visual subgroup of (W, S) , then $(\langle A \rangle, A)$ is also a Coxeter system by Theorem 2 (i), p. 20 of Bourbaki [3].

When studying a Coxeter system (W, S) with Coxeter matrix M it is helpful to have a visual representation of (W, S) . There are two graphical ways of representing (W, S) and we shall use both depending on our needs.

The *Coxeter diagram* (*C-diagram*) of (W, S) is the labeled undirected graph $\Delta = \Delta(W, S)$ with vertices S and edges

$$\{(s, t) : s, t \in S \text{ and } m(s, t) > 2\}$$

such that an edge (s, t) is labeled by $m(s, t)$. Coxeter diagrams are useful for visually representing finite Coxeter groups. If $A \subset S$, then $\Delta(\langle A \rangle, A)$ is the subdiagram of $\Delta(W, S)$ induced by A .

A Coxeter system (W, S) is said to be *irreducible* if its C-diagram Δ is connected. A visual subgroup $\langle A \rangle$ of (W, S) is said to be *irreducible* if $(\langle A \rangle, A)$ is irreducible. A subset A of S is said to be *irreducible* if $\langle A \rangle$ is irreducible.

A subset A of S is said to be a *component* of S if A is a maximal irreducible subset of S or equivalently if $\Delta(\langle A \rangle, A)$ is a connected component of $\Delta(W, S)$. The connected components of the $\Delta(W, S)$ represent the factors of a direct product decomposition of W .

The *presentation diagram* (P-diagram) of (W, S) is the labeled undirected graph $\Gamma = \Gamma(W, S)$ with vertices S and edges

$$\{(s, t) : s, t \in S \text{ and } m(s, t) < \infty\}$$

such that an edge (s, t) is labeled by $m(s, t)$. Presentation diagrams are useful for visually representing infinite Coxeter groups. If $A \subset S$, then $\Gamma(\langle A \rangle, A)$ is the subdiagram of $\Gamma(W, S)$ induced by A . The connected components of $\Gamma(W, S)$ represent the factors of a free product decomposition of W .

For example, consider the Coxeter group W generated by the four reflections in the sides of a rectangle in E^2 . The C-diagram of (W, S) is the disjoint union of two edges labeled by ∞ while the P-diagram of W is a square with edge labels 2.

Let (W, S) and (W', S') be Coxeter systems with P-diagrams Γ and Γ' , respectively. An *isomorphism* $\phi : (W, S) \rightarrow (W', S')$ of Coxeter systems is an isomorphism $\phi : W \rightarrow W'$ such that $\phi(S) = S'$. An *isomorphism* $\psi : \Gamma \rightarrow \Gamma'$ of P-diagrams is a bijection from S to S' that preserves edges and their labels. Note that $(W, S) \cong (W', S')$ if and only if $\Gamma \cong \Gamma'$.

We shall use Coxeter's notation on p. 297 of [5] for the irreducible spherical Coxeter simplex reflection groups except that we denote the dihedral group \mathbf{D}_2^k by $\mathbf{D}_2(k)$. Subscripts denote the rank of a Coxeter system in Coxeter's notation. Coxeter's notation partly agrees with but differs from Bourbaki's notation on p.193 of [3].

Coxeter [4] proved that every finite irreducible Coxeter system is isomorphic to exactly one of the Coxeter systems \mathbf{A}_n , $n \geq 1$, \mathbf{B}_n , $n \geq 4$, \mathbf{C}_n , $n \geq 2$, $\mathbf{D}_2(k)$, $k \geq 5$, \mathbf{E}_6 , \mathbf{E}_7 , \mathbf{E}_8 , \mathbf{F}_4 , \mathbf{G}_3 , \mathbf{G}_4 . For notational convenience, we define $\mathbf{B}_3 = \mathbf{A}_3$, $\mathbf{D}_2(3) = \mathbf{A}_2$, and $\mathbf{D}_2(4) = \mathbf{C}_2$

The *type* of a finite irreducible Coxeter system (W, S) is the isomorphism type of (W, S) represented by one of the systems \mathbf{A}_n , \mathbf{B}_n , \mathbf{C}_n , $\mathbf{D}_2(k)$, \mathbf{E}_6 , \mathbf{E}_7 , \mathbf{E}_8 , \mathbf{F}_4 , \mathbf{G}_3 , \mathbf{G}_4 . The *type* of an irreducible subset A of S is the type of $(\langle A \rangle, A)$.

The C-diagram of \mathbf{A}_n is a linear diagram with n vertices and all edge labels 3. The C-diagram of \mathbf{B}_n is a Y-shaped diagram with n vertices and all edge labels 3 and two short arms of consisting of single edges. The C-diagram of \mathbf{C}_n is a linear diagram with n vertices and all edge labels 3 except for the last edge labelled 4. The C-diagram of $\mathbf{D}_2(k)$ is a single edge with label k . The C-diagrams of $\mathbf{E}_6, \mathbf{E}_7, \mathbf{E}_8$ are star shaped with three arms and all edge labels 3. One arm has length one and another has length two. The C-diagram of \mathbf{F}_4 is a linear diagram with edge labels 3, 4, 3 in that order. The C-diagram of \mathbf{G}_3 is a linear diagram with edge labels 3, 5. The C-diagram of \mathbf{G}_4 is a linear diagram with edge labels 3, 3, 5 in that order.

3 Elementary Quotient Operations

In this section we describe two types of elementary edge quotient operations on a Coxeter system (W, S) of finite rank. The first we call edge label reduction and the second we call edge elimination.

Suppose s and t are distinct elements of S with $m(s, t) < \infty$. Let d be a positive divisor of $m = m(s, t)$, with $d < m$, and let N be the normal closure of the element $(st)^d$ of W . Then a presentation for W/N is obtained from the Coxeter presentation for (W, S) by adding the relator $(st)^d$. As $m = (m/d)d$, the relator $(st)^m$ is derivable from the relator $(st)^d$ and so the relator $(st)^m$ can be removed from the presentation for W/N .

Assume $d > 1$. Then the presentation for W/N is a Coxeter presentation whose P-diagram is obtained from the P-diagram for (W, S) by replacing the label m on the edge (s, t) with the label d . We call the operation of passing from the Coxeter system (W, S) to the quotient Coxeter system $(W/N, \{sN : s \in S\})$ the (s, t) *edge label reduction* from m to d . For example, if we reduce the 4 edge of \mathbf{F}_4 to 2, we obtain the Coxeter system $\mathbf{A}_2 \times \mathbf{A}_2$.

Now assume $d = 1$. We delete from the presentation for W/N the generator t and the relator st and replace all occurrences of t in the remaining relators by s . Suppose r is in S and $k = m(r, s) < \infty$ and $\ell = m(r, t) < \infty$. Then we have the relators $(rs)^k$ and $(rs)^\ell$ in the presentation for W/N . Let d be the greatest common divisor of k and ℓ . Then there are integers a and b such that $d = ak + b\ell$. This implies that $(rs)^d$ is derivable from $(rs)^k$ and $(rs)^\ell$ and so we may add the relator $(rs)^d$ to the presentation for W/N . Then $(rs)^k$ and $(rs)^\ell$ are derivable from $(rs)^d$ and so we can eliminate the relators $(rs)^k$ and $(rs)^\ell$ from the presentation for W/N . We do this for each r in S

such that $m(r, s) < \infty$ and $m(r, t) < \infty$. On the P -diagram level, we have eliminated the edge (s, t) and identified the vertices s and t and we have coalesced each edge (r, s) with label $k < \infty$ with the edge (r, t) with label $\ell < \infty$ to form an edge with label d the greatest common divisor of k and ℓ . If each common divisor d is greater than one, we obtain a Coxeter presentation for W/N . If some common divisor d is one, we delete the corresponding generator r and repeating the above reduction procedure on the presentation of W/N . As the set S of generators is finite, we will eventually stop deleting generators and obtain a Coxeter presentation for W/N with generators the subset S' of $\{sN : s \in S\}$ corresponding to the undeleted elements of S . We call the operation of passing from the Coxeter system (W, S) to the quotient Coxeter system $(W/N, S')$ the (s, t) *edge elimination*. For example, if we eliminate the 3 edge from \mathbf{C}_3 , we obtain the Coxeter system $\mathbf{A}_1 \times \mathbf{A}_1$.

4 The Even Part Isomorphism Invariant

Let (W, S) be a Coxeter system of finite rank. For each pair of elements s, t of S with $m(s, t) < \infty$, let $e(s, t)$ be the even part of $m(s, t)$, that is, $e(s, t)$ is the largest power of 2 that divides $m(s, t)$. If $m(s, t) = \infty$, we define $e(s, t) = \infty$. Let N_e be the normal closure in W of all the elements of the form $(st)^{e(s, t)}$ with $e(s, t) < \infty$, and let $W_e = W/N_e$. Let $\eta : W \rightarrow W_e$ be the quotient homomorphism, and let $S_e = \eta(S)$.

Proposition 4.1 *The pair (W_e, S_e) is a Coxeter system such that if s and t are in S , then $\eta(s) = \eta(t)$ if and only if s and t are conjugate in W . If s and t are nonconjugate elements of S , then the order of $\eta(s)\eta(t)$ is the minimum of the set of all $e(u, v)$ such that u and v are in S and u is conjugate to s and v is conjugate to t . In particular, the order of $\eta(s)\eta(t)$ is a power of 2 or ∞ .*

Proof: The system (W_e, S_e) can be obtained from (W, S) by a sequence of elementary quotient operations. First we can do a series of edge label reductions of all the even labelled edges of the P -diagram of (W, S) to their even parts. Then we do a series of edge eliminations of all the odd labelled edges. Each element of the form $(st)^{e(s, t)}$ with $e(s, t) < \infty$ is in the commutator subgroup of W . Therefore abelianizing W factors through the quotient W/N_e , and so $\eta(s) = \eta(t)$ if and only if s and t are the same odd component

of the P-diagram of (W, S) . Hence $\eta(s) = \eta(t)$ if and only if s and t are conjugate in W by Prop. 3, p. 12 of Bourbaki [3].

Suppose s and t are nonconjugate elements of S and u and v are in S , with $m(u, v) < \infty$, and u is conjugate to s and v is conjugate to t . Then u and v are not conjugate, and so $m(u, v)$ is even, and therefore $e(u, v)$ is a power of 2 greater than 1. In the coalescence of two such edges, the greatest common divisor is the minimum of the two edge labels. Therefore the order of $\eta(s)\eta(t)$ is the minimum of the set of all $e(u, v)$ such that u and v are in S and u is conjugate to s and v is conjugate to t . \square

Theorem 4.2 *Let (W, S) be a Coxeter system of finite rank. For each pair of elements s, t of S with $m(s, t) < \infty$, let $e(s, t)$ be the largest power of 2 that divides $m(s, t)$. Let N_e be the normal closure in W of all the elements of the form $(st)^{e(s,t)}$ with $m(s, t) < \infty$. Then N_e is the normal closure in W of the set of all elements of W of odd order. Therefore N_e is a characteristic subgroup of W that does not depend on the choice of Coxeter generators S .*

Proof: Every element of the form $(st)^{e(s,t)}$ with $m(s, t) < \infty$ has odd order, and so N_e is contained in the normal closure of all the elements of odd order. Let w be an element of odd order, then $\eta(w)$ has odd order in $W_e = W/N_e$. By the previous proposition, (W_e, S_e) is a Coxeter system with all edge labels a power of 2. Therefore $\eta(w)$ is conjugate to an element of odd order of a finite visual subgroup of (W_e, S_e) by [3], Ch. V, §4, Ex. 2. The finite visual subgroups of (W_e, S_e) are direct products of groups of type \mathbf{A}_1 and \mathbf{C}_2 , and so are 2-groups. Therefore W_e has no nontrivial elements of odd order. Hence $\eta(w) = 1$, and so w is in N_e . Thus N_e is the normal closure of all the elements of W of odd order. \square

P. Bahls proved in his Ph.D. thesis [1] that any finitely generated Coxeter group has at most one P-diagram, up to isomorphism, with all edge labels even; see Theorem 5.2 in Bahls and Mihalik [2]. Therefore the isomorphism type of the P-diagram of (W_e, S_e) is an isomorphism invariant of W by Theorem 4.2. We call the isomorphism type of the P-diagram of (W_e, S_e) the *even part isomorphism invariant* of W .

In Figure 1, we illustrate two P-diagrams and their even part isomorphism invariant P-diagrams below them. The even diagrams are not isomorphic, and so the top two P-diagrams represent nonisomorphic Coxeter groups.




Figure 1

5 Matching Theorems

Let (W, S) be a Coxeter system. A *basic subgroup* of (W, S) is a noncyclic, maximal, finite, irreducible, visual subgroup of (W, S) . A *base* of (W, S) is a subset B of S such that $\langle B \rangle$ is a basic subgroup of (W, S) . The theorems in this section are proved in our paper [8].

Theorem 5.1 (Basic Matching Theorem) *Let W be a finitely generated Coxeter group with two sets of Coxeter generators S and S' . Let B be a base of (W, S) . Then there is a unique irreducible subset B' of S' such that $[\langle B \rangle, \langle B \rangle]$ is conjugate to $[\langle B' \rangle, \langle B' \rangle]$ in W . Moreover,*

1. *The set B' is a base of (W, S') ,*
2. *If $|\langle B \rangle| = |\langle B' \rangle|$, then B and B' have the same type and there is an isomorphism $\phi : \langle B \rangle \rightarrow \langle B' \rangle$ that restricts to conjugation on $[\langle B \rangle, \langle B \rangle]$ by an element of W .*
3. *If $|\langle B \rangle| < |\langle B' \rangle|$, then either B has type \mathbf{B}_{2q+1} and B' has type \mathbf{C}_{2q+1} for some $q \geq 1$ or B has type $\mathbf{D}_2(2q+1)$ and B' has type $\mathbf{D}_2(4q+2)$ for some $q \geq 1$. Moreover, there is a monomorphism $\phi : \langle B \rangle \rightarrow \langle B' \rangle$ that restricts to conjugation on $[\langle B \rangle, \langle B \rangle]$ by an element of W .*

Let W be a finitely generated Coxeter group with two sets of Coxeter generators S and S' . A basic subgroup $\langle B \rangle$ of (W, S) is said to *match* a basic subgroup $\langle B' \rangle$ of (W, S') if $[\langle B \rangle, \langle B \rangle]$ is conjugate to $[\langle B' \rangle, \langle B' \rangle]$ in W . A base B of (W, S) is said to *match* a base B' of (W, S') if $\langle B \rangle$ matches $\langle B' \rangle$.

Theorem 5.2 *Let (W, S) be a Coxeter system of finite rank. Let B be a base of (W, S) of type \mathbf{C}_{2q+1} for some $q \geq 1$, and let a, b, c be the elements of B such that $m(a, b) = 4$ and $m(b, c) = 3$. Then W has a set of Coxeter generators S' such that B matches a base B' of (W, S') of type \mathbf{B}_{2q+1} if and only if $m(s, t) = 2$ for all $(s, t) \in (S - B) \times B$ such that $m(s, a) < \infty$.*

Theorem 5.3 *Let B be a base of (W, S) of type \mathbf{C}_{2q+1} for some $q \geq 1$, and let a, b, c be the elements of B such that $m(a, b) = 4$ and $m(b, c) = 3$. Suppose that $m(s, t) = 2$ for all $(s, t) \in (S - B) \times B$ such that $m(s, a) < \infty$. Let $d = aba$, and let z be the longest element of $\langle B \rangle$. Let $S' = (S - \{a\}) \cup \{d, z\}$ and $B' = (B - \{a\}) \cup \{d\}$. Then S' is a set of Coxeter generators for W such that*

1. *The set B' is a base of (W, S') of type \mathbf{B}_{2q+1} that matches B ,*
2. *$m(z, t) = 2$ for all $t \in B'$,*
3. *If $(s, t) \in (S - B) \times \{d, z\}$, then $m(s, t) < \infty$ if and only if $m(s, a) < \infty$, moreover if $m(s, t) < \infty$, then $m(s, t) = 2$.*

Theorem 5.4 *Let (W, S) be a Coxeter system of finite rank, and let $B = \{a, b\}$ be a base of (W, S) of type $\mathbf{D}_2(4q + 2)$ for some $q \geq 1$. Then W has a set of Coxeter generators S' such that B matches a base B' of (W, S') of type $\mathbf{D}_2(2q + 1)$ if and only if either $v = a$ or $v = b$ has the property that if $s \in S - B$ and $m(s, v) < \infty$, then $m(s, a) = m(s, b) = 2$.*

Theorem 5.5 *Let $B = \{a, b\}$ be a base of (W, S) of type $\mathbf{D}_2(4q + 2)$ for some $q \geq 1$. Suppose that if $s \in S - B$ and $m(s, a) < \infty$, then $m(s, a) = m(s, b) = 2$. Let $c = aba$ and let z be the longest element of $\langle B \rangle$. Let $S' = (S - \{a\}) \cup \{c, z\}$ and $B' = \{b, c\}$. Then S' is a set of Coxeter generators of W such that*

1. *The set B' is a base of (W, S') of type $\mathbf{D}_2(2q + 1)$ that matches B ,*
2. *$m(z, b) = m(z, c) = 2$,*
3. *if $(s, t) \in (S - B) \times \{c, z\}$, then $m(s, t) < \infty$ if and only if $m(s, a) < \infty$, moreover if $m(s, t) < \infty$, then $m(s, t) = 2$.*

6 The Even Isomorphism Invariant

Let (W, S) be a Coxeter system of finite rank. A pair of elements (a, b) of S is said to be *special* if (a, b) satisfy the conditions of Theorem 5.4, that is, if $m(a, b) \equiv 2 \pmod{4}$ and either $v = a$ or $v = b$ has the property that if $s \in S - \{a, b\}$ and $m(s, v) < \infty$, then $m(s, a) = m(s, b) = 2$.

Let (a, b) be a pair of elements of S . If (a, b) is special, define $\overline{m}(a, b) = 2$ and if (a, b) is nonspecial, define $\overline{m}(a, b) = m(a, b)$.

Let N_{even} be the normal closure in W of all the elements of the form ab with a and b elements of S such that $m(a, b)$ is odd together with all the elements of the form $(ab)^2$ such that (a, b) is a special pair of elements of S . Let $W_{\text{even}} = W/N_{\text{even}}$, let $\eta : W \rightarrow W_{\text{even}}$ be the quotient homomorphism, and let $S_{\text{even}} = \eta(S)$.

Proposition 6.1 *The pair $(W_{\text{even}}, S_{\text{even}})$ is a Coxeter system such that if s and t are in S , then $\eta(s) = \eta(t)$ if and only if s and t are conjugate in W . If s and t are nonconjugate elements of S , then the order of $\eta(s)\eta(t)$ is the greatest common divisor of the set of all $\overline{m}(u, v)$ such that u and v are in S and u is conjugate to s and v is conjugate to t . In particular, the order of $\eta(s)\eta(t)$ is either even or ∞ .*

Proof: The system $(W_{\text{even}}, S_{\text{even}})$ can be obtained from (W, S) by a sequence of elementary quotient operations. First we reduce to 2 all the edge labels of special edges (a, b) to obtain a Coxeter system with Coxeter matrix $\overline{M} = (\overline{m}(s, t))_{s, t \in S}$. Then we eliminate all the odd labelled edges. Each element of the form either st , with $m(s, t)$ odd, or $(st)^2$, with (s, t) special, is in the commutator subgroup of W . Therefore abelianizing W factors through the quotient W/N_{even} , and so $\eta(s) = \eta(t)$ if and only if s and t are the same odd component of the P-diagram of (W, S) . Hence $\eta(s) = \eta(t)$ if and only if s and t are conjugate in W by Prop. 3, p. 12 of Bourbaki [3].

Suppose s and t are nonconjugate elements of S and u and v are in S , with $\overline{m}(u, v) < \infty$, and u is conjugate to s and v is conjugate to t . Then u and v are not conjugate, and so $\overline{m}(u, v)$ is even. In the coalescence of two such edges, the greatest common divisor is even. Therefore the order of $\eta(s)\eta(t)$ is the greatest common divisor of the set of all $\overline{m}(u, v)$ such that u and v are in S and u is conjugate to s and v is conjugate to t . \square

Let (W, S) be a Coxeter system of finite rank. A base B of (W, S) is said to be of *odd type* if there are elements a and b in B , with $m(a, b)$ odd. A base B of (W, S) is said to be *special* if $B = \{a, b\}$ with (a, b) special.

Theorem 6.2 *Let (W, S) be a Coxeter system of finite rank. Then N_{even} is a characteristic subgroup of W that does not depend on the choice of Coxeter generators S .*

Proof: Observe that N_{even} is the normal closure in W of the commutator subgroups of all the basic subgroups $\langle B \rangle$ of (W, S) such that the base B is either of odd type or special. Let S' be another set of Coxeter generators of W . By the Basic Matching Theorem and Theorem 5.4, the group N_{even} is also the normal closure in W of the commutator subgroups of all the basic subgroups $\langle B' \rangle$ of (W, S') such that the base B' is either of odd type or special. Therefore N_{even} is the normal closure in W of all the elements of the form $s't'$ with s' and t' in S' and $m(s', t')$ odd together with all the elements of the form $(s't')^2$ with (s', t') a special pair of elements of S' . Thus N_{even} is a characteristic subgroup of W that does not depend on the choice of Coxeter generators S . \square

Let (W, S) be a Coxeter system of finite rank. P. Bahls proved in his Ph.D. thesis [1] that any finitely generated Coxeter group has at most one P-diagram, up to isomorphism, with all edge labels even. Therefore the isomorphism type of the P-diagram of $(W_{\text{even}}, S_{\text{even}})$ is an isomorphism invariant of W by Theorem 6.2. We call the isomorphism type of the P-diagram of $(W_{\text{even}}, S_{\text{even}})$ the *even isomorphism invariant* of W . For example, the even isomorphism invariant of the system $\mathbf{D}_2(6)$ is the isomorphism type of the P-diagram of the system $\mathbf{A}_1 \times \mathbf{A}_1$.

7 Basic Characteristic Subgroups

Let (W, S) be a Coxeter system of finite rank. A basic subgroup $\langle B \rangle$ of (W, S) is said to be *stable* if for every set of Coxeter generators S' of W , the base B matches a base B' of (W, S') with $|\langle B \rangle| \leq |\langle B' \rangle|$. Note that a basic subgroup $\langle B \rangle$ of (W, S) is *unstable* precisely when the base B satisfies the conditions of either Theorem 5.2 or 5.4. In particular, every special base is unstable.

Let \mathcal{F} be a family of finite irreducible Coxeter system isomorphism types. Let $N(\mathcal{F})$ be the normal closure in W of the commutator subgroups of all the stable basic subgroups of (W, S) of isomorphism type contained in \mathcal{F} together with the commutator subgroups of all the unstable basic subgroups of (W, S) that match a stable basic subgroup of isomorphism type contained in \mathcal{F} . Let $W(\mathcal{F}) = W/N(\mathcal{F})$, let $\eta : W \rightarrow W(\mathcal{F})$ be the quotient homomorphism, and let $S(\mathcal{F}) = \eta(S)$.

Theorem 7.1 *The pair $(W(\mathcal{F}), S(\mathcal{F}))$ is a Coxeter system that can be obtained from (W, S) by a finite series of elementary edge quotient operations. The group $N(\mathcal{F})$ is a characteristic subgroup of W that does not depend on the choice of Coxeter generators S .*

Proof: Quotienting out the commutator subgroup of a basic subgroup $\langle B \rangle$ of (W, S) can be realized by reducing all the even labelled edges of the C-diagram of $(\langle B \rangle, B)$ to 2 and eliminating all the odd labelled edges of the C-diagram. Therefore $(W(\mathcal{F}), S(\mathcal{F}))$ is a Coxeter system that can be obtained from (W, S) by a finite series of elementary edge quotient operations.

Let S' be another set of Coxeter generators of W . By the Basic Matching Theorem, $N(\mathcal{F})$ is also the normal closure in W of the commutator subgroups of all the stable basic subgroups of (W, S') of isomorphism type contained in \mathcal{F} together with the commutator subgroups of all the unstable basic subgroups of (W, S) that match a stable basic subgroup of isomorphism type contained in \mathcal{F} . Thus $N(\mathcal{F})$ is a characteristic subgroup of W that does not depend on the choice of Coxeter generators S . \square

We call a subgroup of W of the form $N(\mathcal{F})$ a *basic characteristic subgroup*.

Corollary 7.2 *If W is a finitely generated Coxeter group and $N(\mathcal{F})$ is a basic characteristic subgroup of W , then the isomorphism type of $W(\mathcal{F}) = W/N(\mathcal{F})$ is an isomorphism invariant of W .*

8 The Spherical Rank Two Invariant

B. Mühlherr [7] has announced a solution of the isomorphism problem for finitely generated Coxeter groups W such that W has no basic subgroups of rank greater than 2 with respect to some set of Coxeter generators. By the Basic Matching Theorem, if W has no basic subgroups of rank greater

than 2 with respect to some set of Coxeter generators, then W has no basic subgroups of rank greater than 2 with respect to every set of Coxeter generators. Therefore it makes sense to say that W has no basic subgroups of rank greater than 2 without regard to a set of Coxeter generators.

In this section we describe a characteristic subgroup N_2 of a finitely generated Coxeter group W such that $W_2 = W/N_2$ is a Coxeter group with no basic subgroups of rank greater than 2 and such that the isomorphism type of W_2 is an isomorphism invariant of W .

Let \mathbf{X}_n be one of the finite irreducible Coxeter systems $\mathbf{A}_n, \mathbf{B}_n, \mathbf{C}_n, \mathbf{E}_6, \mathbf{E}_7, \mathbf{E}_8, \mathbf{F}_4, \mathbf{G}_3, \mathbf{G}_4$ of rank $n \geq 3$. We now define a characteristic subgroup $N(\mathbf{X}_n)$ of \mathbf{X}_n for each \mathbf{X}_n . Let $N(\mathbf{X}_n)$ be the commutator subgroup of the Coxeter group \mathbf{X}_n if $n \geq 5$ or if $\mathbf{X}_n = \mathbf{A}_4, \mathbf{G}_3$, or \mathbf{G}_4 .

Let a_1, a_2, a_3 be the Coxeter generators of \mathbf{A}_3 indexed so that $m(a_1, a_2) = m(a_2, a_3) = 3$. Let $N(\mathbf{A}_3)$ be the normal closure in the group \mathbf{A}_3 of the element a_1a_3 . Then $N(\mathbf{A}_3)$ is a characteristic subgroup of \mathbf{A}_3 characterized by the property that $N(\mathbf{A}_3)$ is the unique normal subgroup of \mathbf{A}_3 such that $\mathbf{A}_3/N(\mathbf{A}_3)$ is isomorphic to \mathbf{A}_2 according to Table 3 of Maxwell [6].

Let b_1, b_2, b_3, b_4 be the Coxeter generators of \mathbf{B}_4 indexed so that

$$m(b_1, b_4) = m(b_2, b_4) = m(b_3, b_4) = 3.$$

Let $N(\mathbf{B}_4)$ be the normal closure in the group \mathbf{B}_4 of the elements b_1b_2 and b_2b_3 . Then $N(\mathbf{B}_4)$ is a characteristic subgroup of \mathbf{B}_4 characterized by the property that $N(\mathbf{B}_4)$ is the unique normal subgroup of \mathbf{B}_4 such that $\mathbf{B}_4/N(\mathbf{B}_4)$ is isomorphic to \mathbf{A}_2 according to Table 3 of Maxwell [6].

Let c_1, c_2, c_3 be the Coxeter generators of \mathbf{C}_3 such that $m(c_1, c_2) = 3$ and $m(c_2, c_3) = 4$. Let $N(\mathbf{C}_3)$ be the normal closure in the group \mathbf{C}_3 of the element $(c_2c_3)^2$. Then $N(\mathbf{C}_3)$ is a characteristic subgroup of \mathbf{C}_3 characterized by the property that $N(\mathbf{C}_3)$ is the unique normal subgroup of \mathbf{C}_3 such that $\mathbf{C}_3/N(\mathbf{C}_3)$ is isomorphic to $\mathbf{A}_2 \times \mathbf{A}_1$ according to Table 3 of Maxwell [6].

Let c_1, c_2, c_3, c_4 be the Coxeter generators of \mathbf{C}_4 indexed so that

$$m(c_1, c_2) = m(c_2, c_3) = 3 \quad \text{and} \quad m(c_3, c_4) = 4.$$

Let $N(\mathbf{C}_4)$ be the normal closure in the group \mathbf{C}_4 of the element c_1c_3 . Then $N(\mathbf{C}_4)$ is a characteristic subgroup of \mathbf{C}_4 characterized by the property that $N(\mathbf{C}_4)$ is the unique normal subgroup of \mathbf{C}_4 such that $\mathbf{C}_4/N(\mathbf{C}_4)$ is isomorphic to $\mathbf{A}_2 \times \mathbf{A}_1$ according to Table 3 of Maxwell [6].

Let f_1, f_2, f_3, f_4 be the Coxeter generators of \mathbf{F}_4 indexed so that

$$m(f_1, f_2) = m(f_3, f_4) = 3 \quad \text{and} \quad m(f_2, f_3) = 4.$$

Let $N(\mathbf{F}_4)$ be the normal closure in the group \mathbf{F}_4 of the element $(f_2 f_3)^2$. Then $N(\mathbf{F}_4)$ is a characteristic subgroup of \mathbf{F}_4 characterized by the property that $N(\mathbf{F}_4)$ is the unique normal subgroup of \mathbf{F}_4 such that $\mathbf{F}_4/N(\mathbf{F}_4)$ is isomorphic to $\mathbf{A}_2 \times \mathbf{A}_2$ according to Table 3 of Maxwell [6].

Let (W, S) be a Coxeter system of finite rank. Let $N(W)$ be the normal closure in W of the subgroups $N(\langle B \rangle)$ defined above for every base B of (W, S) of rank greater than 2. Let $W^{(2)} = W/N(W)$. Let $\eta : W \rightarrow W^{(2)}$ be the quotient homomorphism, and let $S^{(2)} = \eta(S)$.

Theorem 8.1 *The pair $(W^{(2)}, S^{(2)})$ is a Coxeter system that can be obtained from (W, S) by a finite series of elementary edge quotient operations. The group $N(W)$ is a characteristic subgroup of W that does not depend on the choice of Coxeter generators S .*

Proof: Quotienting out the group $N(\langle B \rangle)$ for each base B of (W, S) of rank greater than 2 can be realized by elementary edge quotient operations. Therefore $(W^{(2)}, S^{(2)})$ is a Coxeter system that can be obtained from (W, S) by a finite series of elementary edge quotient operations.

Let S' be another set of Coxeter generators of W . By the Basic Matching Theorem and the characteristic properties of the groups $N(\mathbf{X}_n)$, the group $N(W)$ defined in terms of the generators S is the same as the group $N(W)$ defined in terms of the generators S' . Thus $N(\mathcal{F})$ is a characteristic subgroup of W that does not depend on the choice of generators S . \square

Corollary 8.2 *If W is a finitely generated Coxeter group, then the isomorphism type of $W^{(2)} = W/N(W)$ is an isomorphism invariant of W .*

It may happen that $(W^{(2)}, S^{(2)})$ has a base of rank greater than 2. To get a quotient system with no bases of rank greater than 2, we may have to quotient out $N(W^{(2)})$, and then perhaps repeat the above quotienting process several times. This leads to a finite nested sequence

$$\{1\} = N^{(1)}(W) \subset N^{(2)}(W) \subset \cdots \subset N^{(\ell)}(W)$$

of characteristic subgroups of W such that if $W^{(i)} = W/N^{(i)}(W)$ and if $\eta_i : W \rightarrow W^{(i)}$ is the quotient homomorphism, then

$$N^{(i+1)}(W) = \eta_i^{-1}(N(W^{(i)}))$$

for each $i = 1, \dots, \ell - 1$, and $W^{(\ell)}$ has no basic subgroups of rank greater than 2, and ℓ is as small as possible. We have that $W^{(i+1)} = (W^{(i)})^{(2)}$ for each $i = 1, \dots, \ell - 1$. Therefore the isomorphism type of $W^{(i)}$ for each $i = 1, \dots, \ell$ is an isomorphism invariant of W . It follows from the Basic Matching Theorem that ℓ does not depend on a choice of Coxeter generators for W , and so ℓ is an isomorphism invariant of W . We call ℓ the *spherical rank 2 class* of W . We have $\ell \geq 1$ with $\ell = 1$ if and only if W has no basic subgroups of rank greater than 2. Figure 2 shows the P-diagrams of a sequence $W^{(1)}, \dots, W^{(\ell)}$ with $\ell = 4$ for the Coxeter group $W = W^{(1)}$.

Define $N_2 = N^{(\ell)}(W)$. Then N_2 is a characteristic subgroup of W such that $W_2 = W/N_2$ has no basic subgroups of rank greater than 2. The isomorphism type of W_2 is an isomorphism invariant of W which we call the *spherical rank 2 isomorphism invariant* of W .

Let $\eta : W \rightarrow W_2$ be the quotient homomorphism, and let $S_2 = \eta(S)$. Then (W_2, S_2) is a Coxeter system that can be obtained from (W, S) by a finite series of elementary edge quotient operations.




Figure 2

9 Conclusion

Let (W, S) be a Coxeter system of finite rank. In this paper, we have described three characteristic subgroups N_e, N_{even}, N_2 of W each leading to a quotient isomorphism invariant of W . It is interesting to note that

$$N_2 \subset N_{even} \subset N_e,$$

and so the quotient isomorphism invariants corresponding to N_e, N_{even}, N_2 are progressively stronger. The algorithm for finding a P-diagram for the system (W_e, S_e) starting from a P-diagram of (W, S) is computational fast.

The algorithm for finding a P-diagram for the system (W_{even}, S_{even}) is slower since it has to determine the bases of (W, S) of type $\mathbf{D}_2(4q+2)$ that satisfy the conditions of Theorem 5.4; but, this algorithm is only slightly slower since the conditions of Theorem 5.4 are easy to check. The algorithm for finding a P-diagram for the system (W_{even}, S_{even}) would most likely be incorporated into an efficient computer program that determines if two finite rank Coxeter systems have isomorphic groups, since the even isomorphism invariant would usually determine that two random finite rank Coxeter systems have nonisomorphic groups.

The algorithm for finding a P-diagram for the system (W_2, S_2) is the slowest, but it is not much slower, since it only has to find a subdiagram of the P-diagram of (W, S) of type $\mathbf{A}_3, \mathbf{C}_3$ or \mathbf{G}_3 before it performs an edge quotient operation on an edge of the subdiagram. If the subdiagram is of type \mathbf{A}_3 or \mathbf{G}_3 , then the edge with label 2 is eliminated. If the subdiagram is of type \mathbf{C}_3 , then the 4 edge label is reduced to 2. The algorithm then repeats the routine of searching for a subdiagram of type $\mathbf{A}_3, \mathbf{C}_3$ or \mathbf{G}_3 and performing the corresponding edge quotient operation.

The algorithm for finding a P-diagram for the system (W_2, S_2) would most likely be useful in an efficient program that determines if two finite rank Coxeter systems have isomorphic groups, since the solution of the isomorphism problem for finite rank Coxeter systems that have no bases of rank greater than 2 is considerably simpler than any general solution of the isomorphism problem.

References

- [1] P. Bahls, Even rigidity in Coxeter Groups, Ph.D. Thesis, Vanderbilt University, 2002.
- [2] P. Bahls and M. Mihalik, Reflection independence in even Coxeter groups, *Geometriae Dedicata* 110 (2005), 63-80.
- [3] N. Bourbaki, *Groupes et algèbres de Lie*, Chapitres 4, 5, et 6, Hermann, Paris, 1968.
- [4] H.S.M. Coxeter, The complete enumeration of finite groups of the form $R_i^2 = (R_i R_j)^{k_{ij}} = 1$, *J. London Math. Soc.* **10** (1935), 21-25.
- [5] H.S.M. Coxeter, *Regular Polytopes*, Dover, New York, 1973.
- [6] G. Maxwell, The normal subgroups of finite and affine Coxeter groups, *Proc. London Math. Soc.* **76** (1998), 359-382.
- [7] B. Mühlherr, The isomorphism problem for Coxeter groups, Submitted to : Proc. Coxeter Legacy Conference, Toronto (May 2004).
- [8] M. Mihalik, J. Ratcliffe, and S. Tschantz, Matching theorems for systems of finitely generated Coxeter groups, preprint 2006, 49pp.